

Photosensitization: Causes and Testing Options

Photosensitization is a serious skin condition in horses and cattle caused by a hazardous combination of certain plants and ultraviolet (UV) light. Certain plants contain photodynamic agents, which then cause a reaction in the animal's body that leads to ultra-sensitive skin. This condition is specific to lightly or non-pigmented animals or areas of an animal that have less hair. Severe skin damage can result and include ulceration, necrosis, and edema.

Photosensitization is generally classified by the source of the photodynamic agent. The three typical classifications are:

- **Primary Photosensitization (Type I):** Occurs when a photodynamic agent is injected, ingested, or absorbed in the skin. Once an agent enters the systemic circulation and is exposed to UV light, damage to the skin's cell membrane occurs.
- **Aberrant Pigment Metabolism (Type II):** Syndrome in which autogenous pigments are photosensitizing porphyrin agents. This syndrome is a congenital defect. (uroporphyrin I, coproporphyrin I, protoporphyrin III)
- **Secondary (Hepatogenous) Photosensitization (Type III):** Most common form observed in livestock. Occurs when an animal ingests plants that contain phylloerythrin. If the animal has liver damage, phylloerythrin will not be excreted into the animal's bile and will eventually be circulated. When phylloerythrin reaches the skin, it will initiate a phototoxic reaction, causing severe skin burns and sloughing.

Causative Toxic Plants

Primary photosensitization:

1. Bishop's weed (*Ammi majus*) furocoumarin
2. Rainlily (*Cooperia pedunculata*) fungal elaboration
3. Dutchman's breeches (*Thamnosma texana*) psoralens
4. St. John's Wort (Hypericin)
5. Buckwheat (Fagopyrin)
6. Coal tar derivatives such as tetracyclines and polycyclic aromatic hydrocarbons

Hepatogenous photosensitization :

1. Moldy bermudagrass (*Cynodon dactylon*) fungal elaboration
2. Lantana (*Lantana camara*) lantadene A and B
3. Sacahuista, blooms only (*Nolina texana*) saponins
4. Kleingrass (*Panicum coloratum*) saponins
5. Puncturevine (*Tribulus terrestris*) saponins

Clinical Findings and Lesions

Regardless of the cause, classical signs of photosensitization are similar. Classical signs of a photosensitive animal are:

- Immediate discomfort and agitation when exposed to light
- Rubbing or scratching of lightly pigmented or exposed skin areas
- Lesions in exposed areas

Main lesions include skin necrosis and ulceration of exposed areas. Exposed areas include, but are not limited to:

- Face
- Sides of udders
- Underside of tongue
- Lips
- Eyelids
- Ears

Differences in distribution are related to differences in skin pigmentation of breeds and individual animals. Affected skin will be red, oozing fluid, and swollen. In severe cases, large crusts of necrotic, black skin may slough off. Affected animals may also be icteric with yellow discoloration of the skin and mucus membranes. Animals may also experience blindness due to corneal opacity.

Testing Recommendations

TVMDL recommends submitting samples for the following tests. In the case of photosensitization, rumen microscopy is of limited use.

- Chemistry Profile - Ruminant
- Histopathology-Biopsy
- Plant Identification

Send samples to the College Station laboratory:

TVMDL

483 Agronomy Road

College Station, Texas 77843-4471

Information on sample shipment can be found at tvmdl.tamu.edu. Contact the laboratory with questions.

